

WHAT IS CLAIMED IS:

1. A nanotube to which is attached a component linked to a label compound capable of being induced to luminesce.
2. A nanotube according to claim 1, wherein said nanotube is graphitic and said luminescence is electrochemiluminescence.
3. A graphitic nanotube according to claim 1, wherein said component is an enzyme biosensor.
4. A composition for the detection of an analyte of interest present in a sample, which composition comprises:
 - (i) a graphitic nanotube containing a functional group and
 - (ii) an assay-performance-substance linked to said functional group, said assay-performance-substance being capable of binding to the analyte-of-interest.
5. A composition for the detection of an analyte of interest present in a sample, which composition comprises:
 - (i) a graphitic nanotube containing a functional group and
 - (ii) an assay-performance-substance linked to said functional group, said assay-performance-substance being bound to the analyte of interest.
6. A composition according to claim 5, further comprising a second assay-performance-substance bound to the analyte, said second assay-performance-substance being linked to a label compound capable of being induced to luminesce.

7. A composition according to claim 4, wherein said assay-performance-substance contains at least one substance selected from the group consisting of

- (i) added analyte of interest or added analogue of said analyte;
- (ii) a binding partner of said analyte or a binding partner of said analogue; and
- (iii) a reactive component capable of binding with the selected substance.

8. A composition according to claim 6, wherein said assay-performance-substance contains at least one substance selected from the group consisting of

- (i) added analyte of interest or added analogue of said analyte;
- (ii) a binding partner of said analyte or a binding partner of said analogue; and
- (iii) a reactive component capable of binding with the selected substance.

9. A method for performing a binding assay for an analyte of interest present in a sample comprising the steps of:

- (a) forming a composition containing
 - (i) said sample
 - (ii) an assay-performance-substance linked to a label compound capable of being induced to luminesce, and
 - (iii) a plurality of functionalized graphitic nanotubes bound to an assay-performance-substance;
- (b) incubating said composition to form a complex which includes a graphitic nanotube and said label compound;
- (c) collecting said complex in a measurement zone;

(d) inducing the label compound in said complex to luminesce, and

(e) measuring the emitted luminescence to measure the presence of the analyte of interest in the sample.

10. A method as recited in claim 9 wherein said complex is collected on the surface of means for inducing luminescence and measuring said luminescence.

11. A method as recited in claim 10 based upon measurement of electrochemiluminescence wherein said complex is collected at an electrode surface.

12. A method for performing a binding assay for an analyte of interest present in a sample based upon measurement of electrochemiluminescence at an electrode surface comprising the steps:

(a) forming a composition containing

(i) said sample

(ii) an assay-performance-substance linked to a label compound capable of being induced to electrochemiluminesce, and

(iii) a plurality of functionalized graphitic nanotubes bound to an assay-performance-substance;

(b) incubating said composition to form a complex which includes a graphitic nanotube and said label compound;

(c) collecting said complex;

(d) causing said collected complex to come in contact with an electrode surface and inducing the label compound in said complex to luminesce by impressing a voltage on said electrode; and

(e) measuring the emitted luminescence at the electrode surface to measure the presence of the analyte of interest in the sample

13. A method for performing a binding assay for an analyte of interest present in a sample based upon measurement of electrochemiluminescence at an electrode surface comprising the steps:

- (a) forming a composition containing
 - (i) said sample
 - (ii) an assay-performance-substance linked to a label compound capable of being induced to electrochemiluminesce, and
 - (iii) a plurality of magnetically responsive suspended graphitic nanotubes bound to an assay-performance-substance;
- (b) incubating said composition to form a complex which includes a graphitic nanotube and said label compound;
- (c) collecting said complex by imposition of a magnetic field on said graphitic nanotubes;
- (d) causing said collected complex to come in contact with an electrode surface and inducing the label compound in said complex to luminescence by imposing a voltage on said electrode; and
- (e) measuring the emitted luminescence at the electrode surface to measure the presence of the analyte of interest in the sample.

14. A method as recited in claim 13 wherein the imposition of said magnetic field causes said complex to collect at the surface of said electrode.

15. A composition of matter for use as a reagent in a microparticulate-based binding assay comprising functionalized graphitic nanotubes bound to an assay-performance-substance and at least one other component selected from the group consisting of:

- (a) electrolyte;
- (b) label compound containing an ECL moiety;

- (c) analyte of interest or an analog of the analyte of interest;
- (d) a binding partner of the analyte of interest or of its analog;
- (e) a reactive component capable of reacting with (c) or (d);
- (f) a reductant; and
- (g) an electrochemiluminescent-reaction enhancer; provided, however, that no two components contained within any reagent composition are reactive with one another during storage so as to impair their function in the intended assay.

16. A reagent as recited in claim 15 containing magnetically responsive graphitic nanotubes.

17. An assay reagent for an assay based upon a binding reaction and the measurement of an electrochemiluminescent phenomenon comprising:

- (a) an electrolyte;
- (b) a plurality of magnetically responsive functionalized graphitic nanotubes bound to an assay-performance-substance; and
- (c) an assay-performance-substance bound to a chemical moiety having electrochemiluminescent properties.

18. A method for performing an assay for an analyte of interest present in a sample comprising the steps of:

- (a) forming a composition containing
 - (i) said sample, and
 - (ii) a functionalized graphitic nanotube bound to a component which is bound to a label compound capable of being induced to luminesce, wherein said component is a substrate of the analyte of interest;

REFERENCES

1. Beaucage SL, Caruthers MH. Deoxynucleoside phosphoramidites, a new class of key intermediates for deoxypolynucleotide synthesis. Tetrahedron Lett 1982; 22:1859-62.
2. Shibata DK, Arnheim N, Martin JW. Detection of human papilloma virus in paraffin-embedded tissue using the polymerase chain reaction. J Exp Med 1988; 167:225-30.
3. Yanofsky, C. et al. (1981) Nucleic Acids Res. 24, 6647-6668.
4. Updyke TV, Nicolson GL. Immunoaffinity isolation of membrane antigens with biotinylated monoclonal antibodies and streptavidin-agarose. Methods Enzymol 1986; 121:717-25.
5. Cardullo RA, Agrawal S, Flores C, Zamecnik DC, Wolf DE. Detection of nucleic acid hybridization by nonradiative fluorescence resonance energy transfer. Proc Natl Acad Sci 1988; 85:8790-4.
6. Ngo TT. Procedure for activating polymers with primary and or secondary hydroxyl groups. Makromol Chem Macromol Symp 1988; 17:224-39.
7. Coutlee F, Bobo L, Mayur K, Yoken RH, Viscidi RP. Immunodetection of DNA with biotinylated RNA probes: A study of reactivity of a monoclonal antibody to DNA-RNA hybrids. Anal Biochem 1989; 181:96-105.
8. Casadei J, Powell MJ, Kenten JH. Expression and secretion of aequorin as a chimeric antibody using a mammalian expression vector. Proc Natl Acad Sci 1990; 87:2047-51.

9. Molecular cloning, a laboratory manual 2nd Ed
Sambrook, J. Cold Spring Harbor Laboratory New
York.
10. Heney, G. and Orr, G.A. (1981) Anal Biochem.
114, 92-96.
11. Mullis KB, Faloona FA. Specific synthesis of
DNA in vitro via a polymerase-catalyzed chain
reaction. Methods Enzymol 1987; 155:335-50.
12. Lyons J, Janssen JWG, Bartram C, Layton M,
Mufti GJ. Mutation of Ki-ras and N-ras
oncogenes in myelodysplastic syndromes. Blood
1988; 71:1707-12.
13. Saiki RK, Gelfand DH, Stoffel S, Scharf SJ,
Higuchi R, Horn GT, Mullis KB, Erlich HA.
Primer-directed enzymatic amplification of DNA
with a thermostable DNA polymerase. Science
1988; 239:487-91.
14. Yee C, Krishnan-Hewlett I, Baker CC, Schlegel
R, Howley PM. Presence and Expression of Human
Papillomavirus sequences in human cervical
carcinoma cell lines. Am J Pathol 1985;
119:361-6.
15. Reddy EP, Reynolds RK, Santo E, Barbacid M. A
point mutation is responsible for the
acquisition of the transforming properties by
the T24 humanbladder carcinoma oncogene.
Nature 1982; 300:149-52.
16. Marmur, J. (1961) J. Mol. Biol 3, 208.